

Hepatology COVID-19 Registry Week 7 Report

2 new cases added to the report since last week


	Chronic liver disease (N=18)	Post-liver transplantation (N=14)
Gender (%)	Female = 10 (55.5%)	Female = 7 (50%)
	Male = 8 (44.5%)	Male = 7 (50%)
Median Age (yr) at time of COVID diagnosis (range)	5 (0.16-20)	8 (0.5-21)
Etiology of liver disease	Biliary atresia = 6	Biliary atresia = 7
	NAFLD = 4	Acute liver failure = 3
	Autoimmune hepatitis = 3	Metabolic disorders = 1
	Metabolic disorder = 1	Tumor = 1
	Others* = 4	Others** = 2
Highest level of care		
Outpatient	5	9
Hospital floor	8	2
ICU	5	3
Immunosuppression at time	Azathioprine = 3	Tacrolimus = 12
of COVID diagnosis	Steroids =1	Steroids = 6
(*some patients were on multiple	No Immunosuppression = 15	Mycophenolate Mofetil = 6
agents)		Sirolimus = 1
		Cyclosporine =1
Specific Treatment for COVID	Hydroxychloroquine + Azithromycin = 3	Azithromycin = 1
(*some patients were on multiple	IVIG = 3	Hydroxychloroquine = 2
agents)	Steroids = 3	IVIG = 1
	Tocilizumab + Sarilumab = 1	Favipiravir = 1
	Rocilizumab Sarilumab = 1	
	Remdesivir = 2	
	Convalescent Plasma = 1	
	Anticoagulation = 1	
	Anakinra = 1	
	Eculizimab = 1	
Highest respiratory support		
None	12	13
Nasal cannula/CPAP/BiPAP	3	1
Mechanical ventilation	2	0
High Frequency Oscillatory	1	0
ventilation		
Pending information	-	-
Final clinical outcome		
Death	0	0
Recovery	14	12
Still active in clinical course	4	2
Pending information	-	-
Liver related Complications	Ascites = 5	-
	Portal Hypertension Bleeding =1	

*Others CLD: idiopathic cholestasis, GVHD/vanishing bile duct syndrome (Post BMT), IFALD with biliary obstruction, PFIC

** Others PLT: IFALD


Two recipients had multivisceral transplantation

Presenting symptoms at time of diagnosis



*Constitutional symptoms reported: Loss of smell/taste , fever, myalgia, fatigue, sore throat # Other symptoms reported: Malaise, anorexia, headache, rash, Kawasaki disease like presentation(MIS-C)

Change done to Immunosuppressive agent (Post LT)



*Stopped, decreased the dose, others and No changes Commonest agent **stopped**: Mycophenolate Mofetil